

Numerical investigation of the contribution of zonules in the accommodation of the human lens using the Finite Element Method


Dr Kehao Wang (NTU)

Dr-Ing Demetrios T. Venetsanos (CU)

Professor Jian Wang (KU)

Professor Barbara K. Pierscioneck (NTU)

Anatomy of human eye



[http://www.skybrary.aero/index.php/Vision_\(OGHFA_BN\)](http://www.skybrary.aero/index.php/Vision_(OGHFA_BN))

Eye:

- Fibrous layer: cornea and sclera
- Vascular layer: choroid, ciliary body, iris (pupil)
- Nervous layer: retina


Lens:

- The lens is held in place by zonular fibres
- Zonular fibres are attached to lens peripheral zone




<http://www.eyemd.com/>

What is presbyopia



- Sometime after the 40th birthday, it is more difficult to read
- With age, the lens gradually hardens and does not change shape
- Reading glasses with plus-power gives the eye the power it needs to see clearly

Accommodation



© Copyright. 2012. University of Waikato. All Rights Reserved.

[1] Helmholtz, H. Über die Akkommodation des Auges. *A. v. Graefe's Arch. Klin. Ophthalmol* **1**, 1–74 (1855)

[2] Schachar, R.A. Zonular function: a new hypothesis with clinical implications. *Ann. Ophthalmol.* **26**, 36–38 (1994)

Why should we care about presbyopia


- 1.04 billion people globally with presbyopia in 2005
- 517 million had no spectacles or inadequate spectacles (of these, 410 million were prevented from performing near tasks in the way they required)
- Vision impairment from uncorrected presbyopia predominantly exists (94%) in the developing world

Holden BA, Fricke TR, Ho SM, et al. Global Vision Impairment Due to Uncorrected Presbyopia. Arch Ophthalmol. 2008;126(12):1731–1739. doi:10.1001/archophth.126.12.1731

- An estimated 23 million Britons suffer from presbyopia, or age-related long-sightedness

<https://www.telegraph.co.uk/news/health/news/8754607/The-4000-eye-implant-to-replace-reading-glasses.html>

Why should we care about presbyopia




<https://www.statista.com/statistics/491337/eyewear-united-kingdom-uk-market-value/glasses.html>


What can we do about it

FE Analysis: unique tool to analyse the response of eye lens


Single vision Lense




Bifocal Lense



Trifocal Lense



Progressive Lense




Reading glasses




Contact lenses



Corneal inlays (*tiny lenses surgically placed in cornea*)




Conductive Keratoplasty




Refractive lens exchange (RLE): invasive procedure that involves replacing the natural lens of the eye with an artificial **IntraOcular Lens (IOL)**


Geometry of FE lens models



Refractive Index Profile



Optical Image




Solid Model


- ✓ Lens geometry (outer shape & internal cortical layers) can be obtained from optical measurements
- ✓ Both 3-D and axisymmetric Finite Element (FE) models can be developed
- ✓ Present work: axisymmetric / 16yo & 35yo lenses

[1] Bahrami M. et al, *Exp Eye Res*, 124:93-99, 2014.

FE models of eye lens



Burd HJ, Judge SJ, Cross JA. Numerical modelling of the accommodating lens. *Vision Research*. 2002;42:2235–2251.



Belaidi, A. and Pierscioneck B.K. Journal of Vision (2007) 7(11):1, 1–12

Modelling zonular fibres

16-year-old lens:

(a) geometry


(b) FE model with
anchorage regions of
zonular fibres on lens
capsule

35-year-old lens:


(c) geometry

(d) FE model with
anchorage regions of
zonular fibres on lens
capsule


Wang K. et al, Sci Rep, 7, 2017.



Modelling zonular fibres



Bounded zonular fibres




Split zonular fibres

- ✓ Material properties of lens nucleus and cortex: linear elastic, isotropic and homogenous.
- ✓ Poisson's ratio: 0.49 (nearly incompressible material)
- ✓ Nonlinear geometric simulations were performed.

Wang K. et al, Sci Rep, 6, 2016.


Exhaustive search scheme



Optical response of the model

Central Optical Power:
$$\text{COP} = \frac{n_1 - n_2}{r_a} + \frac{n_1 - n_a}{r_p} - \frac{(n_1 - n_a)^2}{r_a r_p n_1}, \quad n_1 = 1.42, \quad n_a = 1.336$$

Jones et al., Vision Res. 45, 2352–2366 (2005).



t : thickness
 r_a : RoC of anterior surface
 n_1 : refractive index of aqueous humour



r_p : RoC of posterior surface
 n_2 : refractive index of lens

Optical response of the model



16-year-old

35-year-old

Single



Multi-layer
one




Discontinuous
stress field:

- ✓ Not a problem when examining the displacement field
- ✓ The displacement field is continuous
- ✓ The deformed shape can be considered as acceptable

Wang K. et al, Sci Rep, 6, 2016.

Fitting to *in-vivo* data

- compared to a 29-year-old lens measured *in vivo*



Dubbelman M. et al, Vis Res, 2005.

Conclusions

- ✓ Equatorial zonule: less significant role on lens shape change than anterior & posterior zonular fibres
- ✓ Zonular angle triplets: those providing results more closely aligned to *in-vivo* data give lower accommodative amplitudes → zonular angles change with age and may account for the accommodative loss
- ✓ Capsular thickness should be taken into account in modelling (it affects optical performance of lens)
- ✓ A linear distribution of the modulus of elasticity could describe more accurately the stiffness tensor throughout the eye lens

References

- Helmholtz Hv. Über die Akkommodation des Auges. A. v. Graefe's Arch. Klin. Ophthalmol. 1855; 1: 1-74.
- Schachar RA. Zonular function: a new hypothesis with clinical implications. Annals of Ophthalmology. 1994; 26: 36-38.
- Fincham E. The Mechanism of Accommodation. Br. J. Ophthalmol. Monograph Suppl. 1937; VIII: 1-80.
- Tscherning M. H. v. Helmholtz et la theorie de l'accommodation. Paris. 1909.
- Rohen JW. Scanning electron microscopic studies of the zonular apparatus in human and monkey eyes. Investigative ophthalmology & visual science. 1979; 18: 133-144.
- Wang K, Venetsanos DT, Wang J, Augousti AT, Pierscioneck BK. The importance of parameter choice in modelling dynamics of the eye lens. Scientific Reports. 2017; 7: 16688.
- Schachar RA. The mechanism of accommodation and presbyopia. Kugler Publications. 2012.
- Pierscioneck BK, Regini JW. The gradient index lens of the eye: an opto-biological synchrony. Progress in retinal and eye research. 2012; 31: 332-349.
- Jones CE, Atchison DA, Meder R, Pope JM. Refractive index distribution and optical properties of the isolated human lens measured using magnetic resonance imaging (MRI). Vision research. 2005; 45: 2352-2366.
- Besner S, Scarcelli G, Pineda R, Yun S. In Vivo Brillouin Analysis of the Aging Crystalline Lens In Vivo Brillouin Analysis of the Aging Human Lens. Investigative ophthalmology & visual science. 2016; 57: 5093-5100.
- Scarcelli G, Yun SH. In vivo Brillouin optical microscopy of the human eye. Optics express. 2012; 20: 9197-9202.
- Pierscioneck B, Bahrami M, Hoshino M, Uesugi K, Regini J, Yagi N. The eye lens: age-related trends and individual variations in refractive index and shape parameters. Oncotarget. 2015; 6: 30532-30544.
- Bahrami M, Hoshino M, Pierscioneck B, Yagi N, Regini J, Uesugi K. Optical properties of the lens: An explanation for the zones of discontinuity. Experimental eye research. 2014; 124: 93-99.
- Augusteyn RC, Jones CE, Pope JM. Age-related development of a refractive index plateau in the human lens: evidence for a distinct nucleus. Clinical and Experimental Optometry. 2008; 91: 296-301.

!! Thank you for your attention !!